arXiv:1304.3012v1 [cond-mat.quant-gas] 10 Apr 2013

Cluster Luttinger liquids of Rydberg-dressed gases in optical lattices

Marco Mattioli,[1] Marcello Dalmonte,[1] Wolfgang Lechner,[1] and Guido Pupillo[2]
[1] *Institute for Quantum Optics and Quantum Information,*
Austrian Academy of Sciences, 6020 Innsbruck, Austria,
Institute for Theoretical Physics, University of Innsbruck, 6020 Innsbruck, Austria
[2] *IPCMS (UMR 7504) and ISIS (UMR 7006), University of Strasbourg and CNRS, Strasbourg, France*

We investigate the zero-temperature phases of hard-core bosonic and fermionic gases confined to one dimension and interacting via a class of finite-range soft-core potentials. Using a combination of analytical and numerical methods, we demonstrate the stabilization of critical quantum liquids with qualitatively new features with respect to the Tomonaga-Luttinger liquid paradigm. These features result from frustration and cluster formation in the corresponding classical ground state. Characteristic signatures of these liquids are accessible in state-of-the-art experimental setups with Rydberg-dressed ground state atoms trapped in optical lattices.

PACS numbers: 71.10.Pm,05.30.Jp,32.80.Ee

Frustration is responsible for a variety of phenomena in many-body physics, such as the emergence of gauge symmetries [1], macroscopic ground state degeneracies in strongly-correlated quantum spin and fermionic models [2] and the glass phases [3]. While these effects pertain to two and three dimensions, in one-dimension (1D) frustration usually leads to gapped states in either spin or fermion systems [1], while gapless quantum systems seem to invariably fall within the universality class of Tomonaga-Luttinger (TL) liquids [4–8]. There, according to Haldane's construction [5], the low-energy behavior is described by a free bosonic theory [9], where the asymptotic decay of correlation functions in the microscopic model can be predicted in terms of just a few effective parameters, such as the Luttinger parameter $K = v\pi\mathcal{C}$, with v and $\mathcal{C}$ the sound velocity and compressibility of the liquid [6]. While this can be proven for certain integrable systems, a fundamental question is whether the TL paradigm persists for any kinds of local or non-local interactions.

Non-local (dipolar, van-der-Waals) interactions can be now realized experimentally with ultracold gases of polar molecules [10, 11], groundstate magnetic [12] and Rydberg-excited atoms [13–16] confined to optical lattices [17, 18]. In particular, in two and three dimensions soft-core interactions in gases of groundstate atoms weakly dressed with a Rydberg state using laser light [19, 20] have been shown theoretically to generate exotic phenomena, such as, e.g., supersolid states of matter [21–27]. Quantum phases of 1D soft-core systems have remained so far largely unexplored.

Here, we show that cluster formation due to soft-core potentials [28, 29] on a 1D lattice provides a natural realization of frustrated systems with infinite groundstate degeneracy. This allows for the stabilization of quantum liquids with anomalous cluster-type correlations not captured by the TL model. We derive an effective field theory describing such states by means of bosonization of effective degrees of freedom on an infinitely degenerate ground state manifold and confirm the departure from the TL picture by means of numerical simulations based on the density-matrix-renormalization-group (DMRG) algorithm [30, 31]. Characteristic features of these anomalous *cluster Luttinger* (cL) liquids, are evident in correlation functions and collective excitations. cL liquids should be realizable in experiments for Rydberg-dressed ground state alkali atoms, where the presence of the optical lattice potential allows one to reach the strongly interacting regime.



Figure 1. Sketches of the classical ground states ($t = 0$) of Eq. (1) in the liquid (a), crystalline (b) and cluster (c) phases, for $r_c = 2, 3$ and 4, respectively and particle density $n = 1/4$. Particles are drawn in the lattice as blue spheres (see text). Insets: numerical DMRG computations of density-density correlations $g_2(x)$ for $V/t = 6$ and $L = 48$ in the corresponding phases. For $r_c = 4$ [panel (c)] particles form two kinds of blocks (or *clusters*) with $N_p = 2$ and $N_p - 1$ particles (A and B in the figure, respectively). In the quantum regime, this results in a cluster-type liquid with correlation functions different from a TL liquid [see Insets].

The relevant microscopic Hamiltonian for hard-core bosons or spinless fermions in 1D geometry reads

$$H = -t\sum_i (b_i^\dagger b_{i+1} + \text{c.c.}) + V\sum_i \sum_{\ell=1}^{r_c} n_i n_{i+\ell}. \qquad (1)$$

Here, $b_i^\dagger (b_i)$ are bosonic/fermionic creation (annihilation) operators at the site i, $n_i = b_i^\dagger b_i$, and t is the tunneling rate on a lattice of spacing a, which, from now on, we set as unit length of distances. In the following, $\rho = n$ ($\rho = 1 - n$) for $n \leq 1/2$ ($n > 1/2$), with n the particle density. As shown below, the *soft-core* model of interactions in Eq. (1) emerges naturally in clouds of cold Rydberg-dressed atoms, where the strength V and range r_c of interactions can be independently tuned using external fields. In the following, we focus on the bosonic case, discussing the fermionic one at the end of the manuscript.

Equation (1) has a rich phase diagram, resulting from the combination of the following features of soft-core interactions: (i) their Fourier transform has negative components, which in classical systems are associated with free-space cluster formation [28]; (ii) the competition of r_c with the critical length $r^\star = 1/\rho - 1$ leads to frustration for $r_c > r^\star$. These effects result in the formation of highly degenerate cluster-type ground states in the classical limit ($t = 0$), as well as in novel phases in the quantum regime ($t > 0$). Note that all correlation functions are here particle-hole symmetric, such that the phases for filling n and $1 - n$ are identical.

Fig. 1 shows sketches of the possible ground states for $n = 1/4$ in the classical limit $t = 0$. These phases are: a liquid ($r_c < r^\star$), a crystal ($r_c = r^\star$), and a cluster state ($r_c > r^\star$). The classical liquid [panel (a)] is given by the summation of all particle configurations with the constraint that inter-particle distances are larger than r_c, while the crystal has a particle every $r_c + 1$ lattice sites [panel (b)]. The ground state in the cluster regime is instead highly degenerate. We find that its structure can be computed analytically using the following procedure: we recognize that (i) for $r_c > r^\star$ particles and holes are grouped into blocks of type A and B consisting of N_p and $N_p - 1$ particles, respectively, followed by a number of r_c empty sites [$N_p = 2$ in Fig. 1(c)]. Other block sizes are energetically unfavored. The total number of blocks M for a system of length L is $M = L(1-\rho)/r_c$. (ii) Any exchange of two blocks does not change the energy of the system. The classical ground state then consists of all permutations of blocks A and B with the constraint that their absolute number is fixed (e.g [$AABAAB$], [$ABAAAB$], ...). In this case, the ground state degeneracy grows exponentially with $d = M!/[(M/3)!(2M/3)!]$.

The formation of the correlated block-structure above in the ground state is well captured by the static structure factor $S(k) = \sum_{\ell,j} e^{ik(\ell-j)} g_2(\ell - j)/L$, with $g_2(\ell - j) = \langle n_\ell n_j \rangle - n^2$ the density-density correlation function. This is shown in Fig. 2(a) for the case $n = 3/4$, $r_c = 4 > r^\star$ and $t = 0$, where $S(k)$ peaks at a k-value $k_c < \pi/2$ (solid black line). Numerical DMRG results in the same figure show that this peak persists for $t \simeq V$, where the ground state is a strongly interacting quantum liquid. This signals a departure from the TL picture, where $k_c = \pi/2$ for any strength of interactions, as discussed below. Key questions are (i) what is the underlying mechanism and (ii) what is the effective theory connecting the classical and quantum regimes.



Figure 2. (a) Static structure factor $S(k)$ at fixed $r_c = 4$ and several V/t for a chain of length $L = 48$. Black solid line: classical case ($t = 0$). (b) $S(k)$ for $V/t = 1.5$ and $r_c = 1$, (black circles), 2 (red squares), 3 (blue diamonds), 4 (green up triangles) and 5 (yellow down triangles).

Low-energy theory: For bosonic and fermionic gapless systems, the low-energy physics is usually well captured by the universality class of the TL liquid. In Haldane's formulation, the operators b_j are phenomenologically described by long-wavelength density and phase fluctuation fields, $\varphi(x)$ and $\vartheta(x)$, satisfying $[\vartheta(y), \nabla\varphi(x)] = i\pi\delta(x - y)$, and the density operator reads

$$n(x) = [n - \nabla\varphi(x)'/\pi] \sum_{p\in\mathbb{Z}} e^{2ip(\pi n x - \varphi'(x))}, \qquad (2)$$

with $\varphi'(x) = \pi n x - \varphi(x)/2$. The discreetness of particle density is captured by the last factor in Eq. (2), which determines the behavior of correlation functions beyond the hydrodynamic approximation, as, for example, $g_2(x) = n^2 + K/(2\pi x^2) + \alpha \cos(2\pi n x)/x^{2K}$. Here, α is a non-universal constant and K is the Luttinger parameter, which embodies the effect of interactions in the low-energy theory. The form of $g_2(x)$ implies peaks (if any) in the static structure factor $S(k)$ at momenta commensurate with density excitations, independently of the form and strength of interactions.

A quantitative example for a lattice case is shown in Fig. 2(b), where DMRG results for $S(k)$ are plotted as a function of k for $V/t = 1.5$ and $n = 3/4$ and several values of r_c (a detailed discussion of numerical results is given below). Because of finite range interactions, a peak at $k = \pi/2$ is formed for $r_c \leq r^\star = 3$, consistent with the TL picture. For $r_c > r^\star$, however, the peak shifts to lower values $k < \pi/2$, incompatible with a TL liquid.

These observations require a modification of the effective theory to incorporate inhomogeneous cluster-type density distributions. In the low-density limit $r_c n \gg 1$

and $n \ll 1$, this is achieved by first defining the density as $n(x) = \sum_{m=1}^{M} f(x_m)\delta(x - x_m)$. Here $f(x_m) \in \{1, 2\}$ labels the cluster size (here, clusters with one and two particles). We then introduce a collective *cluster counting* field $\varphi_{cl}(x)$, so that the density operator can be rewritten as $n(x) = \nabla\varphi_{cl}(x) \sum_{m=1}^{M} f(x_m)\delta[\varphi_{cl}(x) - \pi m]$. The field $\varphi_{cl}(x)$ differs sharply from its counterpart $\varphi(x)$ in the standard TL theory, in that it takes integer values at the position of clusters and not of individual particles, as exemplified in Fig. 3(b). This opens the way to novel collective excitations, as shown below.

After introducing $\varphi'_{cl}(x) = (2\pi n\sigma x - \varphi_{cl}(x)/2)$, with $\sigma = M/N_p$ the cluster density, the corresponding functional dependence of $n(x)$ on $\varphi'_{cl}(x)$ reads

$$n(x) = \left[n - \frac{\sigma}{\pi}\nabla\varphi_{cl}(x)'\right] \sum_{k=-\infty}^{\infty} a_k e^{2ik[\varphi_{cl}(x)' - \pi n\sigma x]}, \quad (3)$$

with a_k constants. Similar to TL liquids, in the critical regime the effective low-energy theory is here expected to be a $c = 1$ conformal field theory (corresponding to a compactified boson model); however, Eq. (3) implies that physical observables will differ from those of TL liquids. As an example, $g_2(x)$ now reads $g_2(x) \simeq n^2 + \alpha_1/x^2 + [\alpha_2 \cos(2\pi n\sigma x)]/x^\gamma$, with α_1, α_2 and γ constants. Here, unlike TL liquids, the relation $\alpha_1 = \gamma/(4\pi)$ is not satisfied, preventing a direct estimate of the decay of correlation functions from spectral properties by means of, e.g., level spectroscopy techniques [32].

The position k_c of peaks in $S(k)$ now depends on the interplay between density and interactions, and is predicted to be $k_c = 2n\pi\sigma = 2\pi(1-\rho)/r_c$, with $k_c < \pi/2$ (see also above). As an example, for $r_c = 4$ and $\rho = 1/4$, $k_c = 3\pi/8$ in perfect agreement with classical results [see Fig. 2(a)]. The small deviation of the $\delta = 2\pi/L$ of the theory from the DMRG results will be subject of future studies. Appreciable differences occur for correlation functions and spectral properties. We note that the procedure above allows one to retain high-momentum features due to interactions in the low-energy theory (here corresponding to cluster formation), and can be in principle applied to other problems where the classical ground state presents infinite degeneracies.

Numerical results: We investigate the phase diagram of Eq. (1) by performing numerical DMRG simulations for a given particle density n as a function of V/t and r_c. In order to avoid boundary effects, which can be significant due to the finite range of interactions, we perform finite-size scaling in systems as large as $L = 64$ sites with periodic boundary conditions, keeping up to 1800 states per block and performing 5 sweeps. This ensures numerical errors of order 10^{-7} or less.

An example for $n = 3/4$ is shown in Fig. 3(a). There, in the considered range of parameters, for sufficiently small interactions, the low-energy phase is shown to be always a TL liquid, independent of r_c. This is clearly seen for, e.g., $r_c = 2 < r^\star$, where the system is equivalent to a XXZ model with next-nearest-neighbor interactions and finite magnetization. For $r_c = r^\star = 3$, instead, the TL liquid turns into an insulating Mott phase with a hole every four sites [blue squares, see also sketch in Fig. 1(b)] via a quantum phase transition of the Berezinskii-Kosterlitz-Thouless (BKT) type around $V/t \simeq 4$, as discussed in Ref. [32]. The most interesting behavior occurs for $r_c/r^\star > 1$ and $V/t \gtrsim 1$, where the TL turns into a cL liquid. Below we analyze in detail this regime, and comment on the region with $V/t \gg 1$ and large $r_c/r^\star \gtrsim 5$ (green triangles) towards the end of the paper.



Figure 3. Left panel: numerical zero-temperature phase diagram of Eq. (1) for density $n = 3/4$. The following phases are indicated: Luttinger liquid (red circles), crystal (blue squares), cluster Luttinger liquid (yellow circles). Green triangles define a phase which shows an anomalous entanglement entropy growth (see main text). Right panel: Configurations allowed for the quantum field $\varphi(x)$: in the TLL scenario (top), $\varphi(x)$ takes integer values at the position of each particle (red circles). In the cluster Luttinger liquid (bottom), φ_{cl} describes the cluster density, and not the particle density. This points to a different nature of collective excitations.

We characterize the phase diagram by focussing on: *i)* $S(k)$; *ii)* the asymptotic decay of g_2 and of the one-body density matrix $B_{ij} = \langle b_i^\dagger b_j \rangle$; *iii)* the excitation spectrum $\Omega(k) = E_k/S(k)$ in the Feynman approximation, with

$$E_k = \frac{t}{L}\left[1 - \cos\left(\frac{2\pi k}{L}\right)\right] \langle 0| \sum_{i=0}^{L-1} (b_i^\dagger b_{i+1} + h.c.)|0\rangle. \quad (4)$$

$\Omega(k)$ is an upper bound to the exact spectrum [33].

According to the discussion above, differences between TL and cL liquids are evident in $S(k)$ and g_2: in particular, in the TL phase $S(k)$ exhibits a peak at $k_c = \pi/2$, commensurate with the particle density, while peaks commensurate with the cluster structure emerge in the cL phase, the exact position depending on r_c. For sufficiently strong interactions $V/t \gtrsim 1$, Figs. 2(a) and (b) demonstrate the crossover from TL to cL liquid increasing strength of interactions at fixed $r_c > r^\star$ and as a function of r_c at fixed V/t, respectively.

Correlation functions such as B_{ij} decay algebraically both in the TL and cL phases, validating the assumption that both underlying field theories are $c = 1$ conformal. However, as illustrated in Fig. 4(a), a strong oscillatory behavior for B_{ij} in the cL phase leads to the emergence of new peaks in the momentum distribution $n(k)$ [Fig. 4(b)]. This effect, which may be observed experimentally, is predicted by the effective theory above and is reminiscent of the deformation of the *Bose surface* observed in frustrated, two-dimensional bosonic models [34]. Finally, as mentioned above, we note that established methods to extract K fail in the cL phase by giving non-consistent estimates in the $r_c > r^\star$ region [32]. This further strengthen the departure from the TL scenario.

The excitation spectrum is analyzed in Figs. 4(c) and (d), where $\Omega(k)$ is plotted as a function of k for several values of r_c with fixed $V/t = 3$, and for increasing V/t with fixed $r_c = 4$, respectively. Panel (a) shows that finite-range interactions with $r_c < r^\star$ induce a roton excitation at $k = \pi/2$, commensurate with density excitations, as expected. This is a precursor of the BKT crystallization at $r_c = r^\star$ and $V/t \gtrsim 4$. For $r_c > r^\star$, however, a new roton-type excitation appears at smaller k. This corresponds to the novel cluster-type excitation. This effect is evident in Fig. 4(d), where $\Omega(k)$ develops cluster-type features at $k \gtrsim \pi/4$ already for weak interactions $V/t = 1$ (red circles), reflecting the structure of the classical ground state.

Finally, in the green region with $V/t \gg 1$ and $r_c/r^\star > 1$ in Fig. 3(a), we observe both (i) correlation functions consistent with a cL liquid and (ii) a significant increase of the entanglement entropy. This may point towards a critical phase with central charge $c > 1$. However, strong finite size effects prevent here any conclusive statements on this regime, which we plan to investigate in a future work.The phase diagram of Fig. 3(a) holds for fermions as well, where the TL phase displays dominant density-wave order; hints on possible physics beyond TL for similar Fermi systems have been reported in Ref. [35].



Figure 4. (a) One body density matrix for $r_c = 4$ with interaction strengths $V/t = 1, 3, 5$. (b) Momentum distribution for the same parameters in (a). (c) Dispersion relation at $V/t = 3$: for $r_c = 2$ ($r_c = 5$) a density roton minimum (cluster-like roton minimum) appear at the corresponding peaks in $S(k)$ [see text and Fig. 2(b)]. (d) Dispersion relation for the same parameters in (a) and (b).



Figure 5. (a) Inset: Laser dressing scheme (see text) and resulting Born-Oppenheimer potential $V_{\rm ryd}(i,j)$ for two Rydberg-dressed atoms (thick red line) compared to the square potential V of Hamiltonian (1) (thin blue line). Panel (a): $g_2(x)$ for several strengths $V_{\rm ryd}/t$ and $r_c = 4$ [symbols as in (b)]. (b) $S(k)$ for several $V_{\rm ryd}/t$ in comparison to $V/t = 6$ (black circles).

Experimental setup and observables. The phases above may be realized using ultracold Rydberg atoms trapped in 1D in an optical lattice. Here we consider ^{85}Rb atoms where each ground state $|g\rangle$ is off-resonantly coupled to an excited Rydberg state $|r\rangle$ via, e.g., a two-photon laser with effective Rabi frequency Ω and red detuning Δ, with $|\Delta| \gg \Omega$, [see Inset of Fig. 5(a)]. We choose $|r\rangle \equiv |43\,S_{1/2}\rangle$, $\Omega/(2\pi) = 1$ MHz and $\Delta/(2\pi) = 10$ MHz (we set the Planck' s constant $h = 2\pi$). The microscopic two-body interaction potential reads $V_{\rm ryd}(i,j) = V_{\rm ryd}\{1 + [(i-j)a/r_c]^6\}^{-1}$, with $V_{\rm ryd}/(2\pi) = \Omega^4/(16\pi\Delta^3) \simeq 125$ Hz. The on-site hard-core constraint is enforced using, e.g., Feshbach resonances. The resulting cut-off radius $r_c = (C_6/2\Delta)^{1/6} \simeq 2.2\mu$m is considerably larger than usual spacings $a \simeq 0.4\,\mu$m. We choose a lattice depth $V_0/(2\pi) \gtrsim 13E_R/(2\pi) = 50$ kHz, such that the longitudinal trapping frequency $\omega_\parallel/(2\pi) = E_R\sqrt{V_0/E_R} \simeq 4.4$ kHz $> \max\{t, V_{\rm ryd}\}$ limits the dynamics to the lowest lattice band. Here, $E_R = \pi^2/(2ma^2)$ is the recoil energy of atoms with mass m, while $t/(2\pi) \lesssim 45$ Hz is the hopping rate in the lattice. Transversal trapping frequencies $\omega_\perp \gg \omega_\parallel$ confine the system to a 1D geometry. The small effective decay rate of the dressed ground state $\gamma_{\rm eff} = (\Omega/2\Delta)^2\gamma_r \simeq 1-10$ Hz ensures dominant Hamiltonian dynamics, with γ_r the bare decay rate of $|r\rangle$. Figure 5 shows that $g_2(x)$ and $S(k)$ using $V_{\rm ryd}(i,j)$ are essentially indistinguishable from those of the potential V of Eq.(1) for $V_{\rm ryd}/t \gg 1$. This opens the way to the observation of the phases above in experiments.

In conclusion, we have shown that soft-core potentials support quantum liquid phases beyond the Tomonaga-Luttinger scenario. These results, which derive from the existence of a macroscopically degenerate ground state in the classical limit, are of immediate interest for experiments with ultracold Rydberg atoms. Interesting extensions of this work include the search for exotic phases in 2D, such as, e.g., frustration-induced Bose metals [36] and emergent gauge theories [1].

Acknowledgements We thank M. Cheneau, F. Cinti, A. Glätzle, C. Gross, A. Läuchli, R. Nath, T. Pohl, P. Zoller for useful discussions, and F. Ortolani for help with the DMRG code. M.M. acknowledges the Marie Curie Initial Training Network COHERENCE for financial support.M.D. acknowledges support by the European Commission via the integrated project AQUTE, and by the Austrian Science Fund FWF (SFB FOQUS F4015-N16).W.L. acknowledges support by the Austrian Science Fund through SFB F40 FOQUS, and by the Institut für Quanteninformation. G.P. acknowledges for hospitality by the KITP in Santa Barbara and support by the European Commission via ERC-St grant "ColdSIM" (grant agreement 307688), EOARD, and UdS via Labex NIE and IdEX.

[1] *Introduction to frustrated magnetism*, edited by C. Lacroix, P. Mendels and F. Mila, Springer (Heidelberg) (2011).
[2] P. A. Lee, N. Nagaosa and X.-G. Wen, Rev. Mod. Phys. **78**, 17 (2006).
[3] H. Shintani and H. Tanaka, Nat. Phys. **2**, 200 (2006).
[4] T. Giamarchi, *Quantum Physics in one dimension*, *Oxford University press* (2003).
[5] F. D. M. Haldane, Phys. Rev. Lett., **47**, 1840 (1981).
[6] A. O. Gogolin, A. A. Nersesyan, A. M. Tsvelik, *Bosonization and strongly correlated systems*, (Cambridge University press, Cambridge, 1998), and references therein.
[7] M. A. Cazalilla, J. Phys. B: At. Mol. Opt. Phys. **37**, 7 (2004).
[8] M. A. Cazalilla, R. Citro, T. Giamarchi, E. Orignac and M. Rigol, Rev. Mod. Phys. **83**, 1405 (2011).
[9] P. Di Francesco, P. Mathieu and D. Sénéchal, *Conformal Field Theory*, Springer-Verlag, New York (1997).
[10] L. D. Carr, D. De Mille, R. V. Krems and J. Ye, New J. Phys. **11**, 055049 (2009).
[11] D. Jin and J. Ye, Chem. Rev. **112**, 4801 (2012).
[12] T. Koch, T. Lahaye, J. Metz, B. Fröhlich, A. Griesmaier, and T. Pfau, Nat. Phys. **4**, 218-222 (2008); G. Bismut, B. Pasquiou, E. Maréchal, P. Pedri, L. Vernac, O. Gorceix and B. Laburthe-Tolra, Phys. Rev. Lett. **105**, 040404 (2010); M. Lu, N. Q. Burdick and B. L. Lev, Phys. Rev. Lett. **108**, 215301 (2012); K. Aikawa, A. Frisch, M. Mark, S. Baier, A. Rietzler, R. Grimm and F. Ferlaino, Phys. Rev. Lett. **108**, 210401 (2012).
[13] D. Comparat and P. Pillet, J. Opt. Soc. Am. B **27**, A208 (2010); R. Löw, H. Weimer, J. Nipper, J. B. Balewski, B. Butscher, H. P. Büchler, T. Pfau, J. Phys. B **45**, 113001 (2012).
[14] J. D. Pritchard, D. Maxwell, A. Gauguet, K. J. Weatherill, M. P. A. Jones and C. S. Adams, Phys. Rev. Lett. **105**, 193603 (2010); M. Viteau, M. G. Bason, J. Radogostowicz, N. Malossi, D. Ciampini, O. Morsch and E. Arimondo, Phys. Rev. Lett. **107**, 060402 (2011); T. Peyronel, O. Firstenberg, Qi-Yu Liang, S. Hofferberth, A. V. Gorshkov, T. Pohl, M. D. Lukin and V. Vuletić, Nature **488**, 57 (2012); P. Schauß, M. Cheneau, M. Endres, T. Fukuhara, S. Hild, A. Omran, T. Pohl, C. Gross, S. Kuhr and I. Bloch, Nature **491**, 87 (2012); M. Viteau, P. Huillery, M. G. Bason, N. Malossi, D. Ciampini, O. Morsch, E. Arimondo, D. Comparat and P. Pillet, Phys. Rev. Lett. **109**, 053002 (2012); M. Robert-de-Saint-Vincent, C. S. Hofmann, H. Schempp, G. Günter, S. Whitlock, M. Weidemüller, Phys. Rev. Lett. **110**, 045004 (2013); T. Baluktsian, B. Huber, R. Löw and T. Pfau, Phys. Rev. Lett. **110**, 123001 (2013).
[15] T. F. Gallagher, *Rydberg Atoms*, Cambridge University Press (1994).
[16] M. Saffman, T. G. Walker and K. Mølmer, Rev. Mod. Phys. **82**, 2313 (2010).
[17] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein and T. Pfau, Rep. Prog. Phys. **72**, 126401 (2009).
[18] M. A. Baranov, M. Dalmonte, G. Pupillo and P. Zoller, Chem. Rev. **112**, 5012 (2012).
[19] L. Santos, G. V. Shlyapnikov, P. Zoller and M. Lewenstein, Phys. Rev. Lett. **85**, 1791 (2001).
[20] G. Pupillo, A. Micheli, M. Boninsegni, I. Lesanovsky and P. Zoller, Phys. Rev. Lett. **104**, 223002 (2010).
[21] L. P. Pitaevskii, Sov. Phys. JETP **13**, 451 (1961); E. P. Gross, J. Math. Phys. **4**, 195 (1963).
[22] Y. Pomeau and S. Rica, Phys. Rev. Lett. **72**, 2426 (1994).
[23] M. Boninsegni and N. V. Prokof'ev, Rev. Mod. Phys. **84**, 759 (2012).
[24] F. Cinti, P. Jain, M. Boninsegni, A. Micheli, P. Zoller and G. Pupillo, Phys. Rev. Lett. **105**, 135301 (2010).
[25] N. Henkel, R. Nath and T. Pohl, Phys. Rev. Lett. **104**, 195302 (2010).
[26] J. Honer, H. Weimer, T. Pfau and H. P. Büchler, Phys. Rev. Lett. **105**, 160404 (2010).
[27] F. Cinti, T. Macrí, W. Lechner, G. Pupillo and T. Pohl, arXiv:1302.4576.
[28] C. N. Likos, A. Lang, M. Watzlawek and H. Löwen, Phys. Rev. E **63**, 031206 (2001).
[29] B. M. Mladek *et al.*, Phys. Rev. Lett. **96**, 045701 (2006); C. N. Likos, B. M. Mladek, D. Gottwald and G. Kahl, J. Chem. Phys. **126**, 224502 (2007); B. M. Mladek, P. Charbonneau and D. Frenkel, Phys. Rev. Lett. **99**, 235702 (2007); D. Coslovich, L. Strauss and G. Kahl, Soft Matter **7**, 2127 (2011).
[30] S. R. White, Phys. Rev. Lett. **69**, (1992) 2863.
[31] U. Schollwöck, Rev. Mod. Phys. **77**, (2005) 259.
[32] See supplemental information.
[33] R. P. Feynman, *Statistical Mechanics*, Advanced Book Program, Westview Press (1998).
[34] C. N. Varney, K. Sun, V. Galitski and M. Rigol, Phys. Rev. Lett. **107**, 077201 (2011).
[35] A. K. Zhuravlev and M. I. Katsnelson, Phys. Rev. B **61**, 534 (2000).
[36] M. S. Block, R. V. Mishmash, R. K. Kaul, D. N. Sheng, O. I. Motrunich and M. P. A. Fisher, Phys. Rev. Lett. **106**, 046402 (2011).